









INDICATORS (%)

Reg. V Phil.

❖ Women with a live birth in the 5 years preceding the survey who received antenatal care (ANC) from:		
✓ doctor	30.0	38.1
✓ nurse/midwife	55.6	49.5
✓ traditional birth attendant/other	8.6	6.5
❖ Women with a live birth in the 5 years preceding the survey who received 2 or more doses of tetanus toxoid injections for their last birth	29.1	37.3
❖ Live births in the 5 years preceding the survey with delivery assistance from:		
✓ doctor	19.5	33.6
✓ nurse	2.1	1.1
✓ midwife	26.2	25.1
✓ hilot	50.2	37.1
❖ Live births in the 5 years preceding the survey that were delivered at:		
✓ home	76.3	61.4
✓ government health facility	16.4	24.2
✓ private health facility	5.5	13.7
❖ Women with a live birth in the 5 years preceding the survey who had first postnatal check-up their last birth at/by:		
✓ home	68.2	45.5
✓ government health facility	25.8	35.1
✓ private health facility/doctor/midwife	6.1	18.8
❖ Children age 12-23 months with no vaccinations at any time preceding the survey	5.6	7.3
❖ Children under 5 years with symptoms of acute respiratory infection (ARI) in the two weeks preceding the survey	9.6	10.2
❖ Children under 5 years with fever in the two weeks preceding the survey	25.6	23.8
❖ Children under 5 years with ARI and/or fever for whom treatment was sought from a health facility/provider	38.2	46.3
❖ Children under 5 years with diarrhea in the two weeks preceding the survey	11.4	10.6
❖ Mothers with births in the 5 years preceding the survey who know about ORS packets	90.2	91.8
❖ Among children born in the 5 years preceding the survey who were ever breastfed, those who started breastfeeding w/ in an hour of birth	34.9	54.0
❖ Among children born in the 3 years preceding the survey, the median duration in months of:		
✓ exclusive breastfeeding	1.9	0.8
✓ predominant breastfeeding	2.4	2.9
✓ any breastfeeding	14.8	14.1
❖ Children age 6-59 months old who received vitamin A supplements in the 6 months preceding the survey	70.9	76.0
❖ Children under 5 years who received iron drops/syrup during the 6 months preceding the survey	53.4	63.3





ABOUT THE SURVEY

The 2003 National Demographic and Health Survey (NDHS) is the eighth in a series of nationwide surveys conducted every five years since 1968.

The survey is designed to provide up-to-date information on population, family planning and health to assist policymakers and program managers in evaluating and designing strategies for improving health and family planning services in the country.

It covered 12,586 households, and completed interviews for 13,633 women 15-49 years old and 4,766 men 15-54 years old.

The 2003 NDHS was implemented by the National Statistics Office (NSO) with technical assistance from ORC Macro and financial support from the United States Agency for International Development (USAID).



For additional information on the results of the 2003 Philippines Demographic and Health Survey, please contact:

In the Philippines:

National Statistics Office
Solicarel Building
Ramon Magsaysay Boulevard
P.O. Box 779
Sta. Mesa, Manila, Philippines
☎ (632) 716 9368/713 7081
☎ (632) 713 7074/714 1715
Internet: <http://www.census.gov.ph>
E-mail: info@census.gov.ph



In the United States:

MEASURE/DHS+
ORC Macro
11785 Beltsville Drive, Suite 300
Calverton, MD 20705 USA
☎ (301) 572 0200
☎ (301) 572 0999
Internet: <http://www.measuredhs.com>
E-mail: reports@orcmacro.com



National Demographic and Health Survey (2003 NDHS)

Fact Sheet



Mayon Volcano

MATERNAL AND CHILD HEALTH and INFANT MORTALITY in BICOL REGION

Philippines